

Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices

UNITED STATES DEPARTMENT OF AGRICULTURE



BULLETIN No. 450



Contribution from the Bureau of Plant Industry
WM. A. TAYLOR, Chief

Washington, D. C.

PROFESSIONAL PAPER

December 4, 1916

IMPROVEMENT OF GHIRKA SPRING WHEAT IN
YIELD AND QUALITY.

By J. ALLEN CLARK,

Scientific Assistant in Western Wheat Investigations, Office of Cereal Investigations.

CONTENTS.

| | Page. | | Page. |
|--|-------|---------------------------------|-------|
| Introduction..... | 1 | Experiments—Continued. | |
| History and description of Ghirka Spring wheat..... | 2 | Summary of yields..... | 8 |
| Experiments..... | 3 | Milling and baking quality..... | 9 |
| Comparative yields..... | 3 | Improvement by selection..... | 12 |
| | | Conclusions..... | 18 |

INTRODUCTION.

A demand for hardier and more drought-resistant wheats was created with the progress of settlement of the drier portions of the Great Plains area. In response to this demand, the United States Department of Agriculture began about 1898 to improve the wheat crop of that area by the introduction from eastern and southern Russia of varieties which were thought to possess hardness and drought resistance. To determine the value of these varieties they were tested at agricultural experiment stations in different localities in the Great Plains area. The principal economic result of this work was the introduction of Kharkof winter wheat and Kubanka durum wheat.¹

Some of the other varieties obtained were found of value for dry-land areas. Among them was the Ghirka Spring wheat, which was both productive and drought resistant, but comparatively low in milling value. Its improvement in yield and quality is the subject of the present paper.

¹ Carleton, M. A. Hard wheats winning their way. *In* U. S. Dept. Agr. Yearbook, 1914, p. 391-420, fig. 22-25, pl. 35-41. 1915.

HISTORY AND DESCRIPTION OF GHIRKA SPRING WHEAT.

The Ghirka Spring is the principal variety of beardless red spring wheat grown in Russia, particularly in southern Russia and the Volga River district. It forms a large part of the wheat exported from Russia.¹ This wheat has been introduced into this country several times. During the period from 1898 to 1904, inclusive, eight lots were obtained by the Office of Cereal Investigations of the United States Department of Agriculture. These lots are recorded as Cereal Investigations Nos. 1046, 1047, 1051, 1192, 1517, 1534, 2644, and 2646.

Other importations of this variety of wheat have been made by Russian immigrants. Joseph Dukart, who settled at New England, N. Dak., brought a 2-pound lot from Russia in 1905. From the



FIG. 1.—Heads of eight varieties of wheat grown at the Dickinson substation: (1) Kubanka durum; (2) Arnautka durum; (3) Preston; (4) Ghirka Spring; (5) Rysting Fife; (6) Marquis; (7) Crossbred Bluestem; and (8) Haynes Bluestem.

increase of this, several thousand acres are now grown annually in western North Dakota as "Russian" wheat. So far, however, the variety has never become commercially important in this country, though its acreage may be expected to increase.

The Ghirka Spring wheat has been placed in the Fife group of spring common wheat by most writers, as its characters are essentially the same as those of the varieties of that group (fig. 1). It differs from the Red Fife varieties in that it is earlier, has pubescent leaves, and the spike is a little more slender and distinctly more tapering at the tip. The kernel is slightly longer, a paler red, and a little softer.

¹ Carleton, M. A. *Triticum vulgare*: Ghirka. In U. S. Dept. Agr., Bur. Plant Indus. Bul. 66 (Seeds and plants imported), no. 6002. 1905.

There is a variety known as Ghirka Winter (C. I. No. 1438), which apparently differs from Ghirka Spring only in the winter habit and in having a shorter, stouter spike.

EXPERIMENTS.

This paper contains the results of experiments with only one of the department's introductions, Ghirka Spring (C. I. No. 1517) from Grodno Province in Russian Poland.

The experimental data are discussed under three separate topics, yield, quality, and improvement. The yield of the original unselected Ghirka Spring wheat is compared with that of three standard varieties at a series of seven stations during a period of seven years. The milling and baking qualities also are compared with those of the same three standard varieties grown at the Dickinson (N. Dak.) sub-station during three different years. Finally, the progress being made in improving both yield and quality by pure-line selection is shown.

COMPARATIVE YIELDS.

Ghirka Spring wheat has been tested in comparison with other domestic and foreign wheats at several agricultural experiment stations in the Great Plains area. Some of these tests have been conducted by the State stations, some by the United States Department of Agriculture, and some by the two in cooperation.

At the time this wheat was included in the varietal tests at the experiment stations in the northern part of the Great Plains, Blue-stem and Fife were the standard wheats grown in that district, while durum wheat was becoming better known and its acreage increasing. The experiment stations were testing several varieties and strains of Fife and Bluestem wheats and many new varieties of durum wheat which were then being imported. The object of these varietal experiments was to determine which group of wheat was best adapted to each locality.

The work at the start was considered a local problem. The varietal tests at each station were practically independent of those at any other. As the work progressed, the best adapted groups and varieties became more and more evident at each testing station. When the results from all the stations in one part of this area are compared only group adaptations usually are shown. The best variety in each group has been not always the same at all stations. In some cases the variety leading at one station has not been grown at some of the other stations or, if grown, has been discarded if the yields were not satisfactory. For these reasons it is difficult to compare the results from individual varieties at a group of stations. It is possible, however, to present yields of Ghirka Spring wheat (C. I. No. 1517) from seven experiment stations in the northern Great Plains during the

period of seven years from 1908 to 1914, inclusive. Yields of Kubanka durum wheat (C. I. No. 1440) and Haynes Bluestem wheat (C. I. No. 2874, Minn. No. 169) for the same years at the same stations are given for purposes of comparison.

It is desirable to compare the performance of Ghirka Spring with that of some other Fife wheat; but no one variety of Fife wheat other than Ghirka has been grown at all of the seven stations during this entire period. However, the Rysting Fife (C. I. No. 3022) has been grown at more stations during the period than any other variety of this group and is chosen for comparison. Yields of Glyndon Fife (C. I. No. 2873) have been substituted at the stations where the Rysting was not grown. The two varieties are very similar in appearance and are only different strains of Fife wheat.

The seven stations for which results ¹ are presented are Moccasin, Mont.; Williston, Dickinson, and Edgeley, N. Dak.; and Brookings, Highmore, and Newell, S. Dak. The location and elevation of these stations are shown in figure 2. At all the stations except Edgeley, the work was conducted cooperatively by the United States Department of Agriculture and the State experiment stations. At Edgeley the tests were conducted entirely by the State,




FIG. 2.—Sketch map of the northern Great Plains area, showing the location and elevation (in feet above sea level) of the seven experiment stations, results from which are discussed in this paper.

and the results given are quoted from the published annual reports of that substation for 1908 to 1913, inclusive. The results for 1914, not yet published, were kindly furnished by O. A. Thompson, superintendent of the Edgeley substation.

The annual and average yields of four varieties of wheat, Kubanka durum, Ghirka Spring, Rysting or Glyndon Fife, and Haynes Bluestem, grown at seven northern Great Plains experiment stations for the 7-year period from 1908 to 1914, inclusive,² are shown in Table I. The average yield of each variety for each station and also the average for all stations are shown graphically in figure 3.

¹ These data have been accumulated at the various stations by the following members of the scientific staff of the Office of Cereal Investigations: Manley Champlin (Brookings, S. Dak.); Charles H. Clark, J. A. Clark, and R. W. Smith (Dickinson, N. Dak.); J. D. Morrison (Highmore, S. Dak.); E. L. Adams and N. C. Donaldson (Moccasin, Mont.); Cecil Salmon and J. H. Martin (Newell, S. Dak.); and F. R. Babcock (Williston, N. Dak.).

² The manuscript of this bulletin was prepared in the spring of 1915, but publication has been unavoidably delayed. This bulletin includes experimental results only to the end of 1914.

The season of 1908 was a favorable one in regard to rainfall at all the stations here considered. Rather conflicting results were obtained with the Ghirka wheat at the different stations. The principal agreement in the tests was the outstanding yield of the Kubanka, surpassing the other varieties of wheat at Williston, Brookings, Highmore, and Newell, and equaling the yields of the Haynes and Ghirka Spring for first

place at Moccasin and Dickinson, respectively. The Ghirka outyielded the other varieties at Edgeley, but, on the other hand, gave the lowest yield at Brookings, Highmore, and Newell. The exceptionally low yield of the Ghirka at Highmore and Brookings was due, in part at least, to rust. The average yield at the seven stations showed Kubanka first, Rysting-Glyndon second, Haynes third, and Ghirka fourth.

The season of 1909 was unusually favorable at all the stations. Kubanka durum wheat gave the highest yields at Moccasin, Williston, Dickinson, Edgeley, and Newell. The Ghirka led at Brookings and Highmore, ranked second at Moccasin, third at Williston, and fourth at Dickinson, Edgeley, and Newell. Under the humid conditions of this year rust again caused a reduction in the yield of the Ghirka. For all stations the Kubanka ranked first, Ghirka second, Haynes third, and Rysting-Glyndon fourth.

The dry season of 1910 reduced the yields at all stations, and the results were entirely different from those obtained in previous years.




FIG. 3.—Diagram showing the average yields of four varieties of spring wheat and the annual and seasonal precipitation at seven experiment stations in the northern Great Plains area for the 7-year period from 1908 to 1914, inclusive.

The Kubanka produced relatively low yields at all stations. This was caused by the sterility of the florets, probably induced by a combination of drought factors.¹ The Ghirka ranked first in yield at Moccasin, Williston, Dickinson, and Newell; second at Edgeley and Brookings; and fourth at Highmore. The results of this year showed that the Ghirka variety possessed real value as a drought-resistant wheat. For all stations the Ghirka ranked first, Rysting-Glyndon second, Haynes third, and Kubanka fourth.

TABLE I.—Yields of the Ghirka Spring and three other varieties of wheat grown at seven experiment stations in the northern Great Plains area, 1908 to 1914, inclusive.

| Station and variety. | C. I. No. | Yield per acre (bushels). | | | | | | | Average. |
|---------------------------------------|-----------|---------------------------|------|------|-------|------|--------|--------|----------|
| | | 1908 | 1909 | 1910 | 1911 | 1912 | 1913 | 1914 | |
| Moccasin, Mont.: | | | | | | | | | |
| Kubanka..... | 1440 | 5.0 | 37.8 | 8.8 | 33.0 | (a) | 30.7 | 23.0 | 23.1 |
| Ghirka..... | 1517 | 3.3 | 34.2 | 13.2 | 25.2 | (a) | 29.0 | 23.0 | 21.3 |
| Rysting..... | 3022 | 2.5 | 29.0 | 13.0 | 27.0 | (a) | 26.7 | 23.0 | 20.2 |
| Haynes..... | 2874 | b 5.0 | 29.7 | 9.1 | 18.7 | (a) | c 22.8 | c 18.6 | 17.7 |
| Williston, N. Dak.: | | | | | | | | | |
| Kubanka..... | 1440 | 12.6 | 39.1 | 11.0 | 8.9 | 51.0 | 33.0 | 53.8 | 29.9 |
| Ghirka..... | 1517 | c 10.7 | 33.2 | 20.2 | 12.1 | 51.7 | 22.2 | 40.4 | 27.2 |
| Glyndon..... | 2873 | 9.8 | 31.2 | 13.9 | 8.0 | 44.7 | 28.2 | 49.2 | 26.4 |
| Haynes..... | 2874 | 10.5 | 34.3 | 8.8 | b 9.2 | 44.3 | 30.0 | 42.5 | 25.8 |
| Dickinson, N. Dak.: | | | | | | | | | |
| Kubanka..... | 1440 | 23.5 | 33.7 | 14.9 | 3.8 | (a) | 26.7 | 14.2 | 19.5 |
| Ghirka..... | 1517 | 23.5 | 28.9 | 28.6 | 9.5 | (a) | 26.6 | 11.3 | 21.4 |
| Rysting..... | 3022 | 22.3 | 33.0 | 20.7 | 7.7 | (a) | 28.1 | 12.4 | 20.7 |
| Haynes..... | 2874 | 21.4 | 30.0 | 13.1 | 8.6 | (a) | 24.8 | 8.3 | 17.7 |
| Edgeley, N. Dak.: ^d | | | | | | | | | |
| Kubanka (Edgeley No. 6)..... | | 11.5 | 27.8 | 5.8 | 2.3 | 34.5 | 25.1 | 17.0 | 17.7 |
| Ghirka (Edgeley No. 162)..... | | 12.4 | 19.7 | 7.9 | 2.0 | 20.1 | 30.6 | 9.0 | 14.5 |
| Rysting (Edgeley No. 3)..... | | 10.1 | 20.3 | 6.2 | 1.6 | 21.7 | 25.2 | 7.0 | 13.2 |
| Haynes (Edgeley No. 46)..... | | 7.9 | 19.9 | 9.0 | 2.0 | 23.5 | 27.8 | 4.0 | 13.4 |
| Brookings, S. Dak.: | | | | | | | | | |
| Kubanka..... | 1440 | 16.1 | 11.8 | 12.7 | 1.2 | 28.0 | 28.3 | 15.0 | 16.2 |
| Ghirka..... | 1517 | 8.8 | 17.7 | 16.2 | .8 | 18.0 | 26.7 | c 8.7 | 13.8 |
| Rysting..... | 3022 | 14.9 | 15.5 | 16.9 | 3.0 | 16.8 | 20.0 | 9.2 | 13.8 |
| Haynes..... | 2874 | 11.6 | 16.9 | 15.7 | 2.8 | 15.8 | 20.2 | 2.5 | 12.2 |
| Highmore, S. Dak.: | | | | | | | | | |
| Kubanka..... | 1440 | 22.7 | 17.0 | 8.0 | 0 | 1.1 | 2.0 | 19.7 | 10.1 |
| Ghirka..... | 1517 | 4.8 | 17.7 | 6.2 | 0 | 2.4 | 6.5 | 11.3 | 7.0 |
| Glyndon..... | 2873 | 13.9 | 15.8 | 12.2 | 0 | 0 | 8.7 | 7.3 | 8.3 |
| Haynes..... | 2874 | 14.7 | 17.2 | 10.0 | 0 | 2.1 | b 7.7 | 7.5 | 8.5 |
| Newell, S. Dak.: | | | | | | | | | |
| Kubanka..... | 1440 | 24.9 | 21.4 | 5.3 | 0 | 0 | 15.6 | c 5.9 | 10.4 |
| Ghirka..... | 1517 | 16.2 | 11.7 | 12.8 | 0 | 1.9 | 16.3 | c 6.1 | 9.3 |
| Rysting..... | 3022 | 19.3 | 15.0 | 10.3 | 0 | 0 | 15.0 | 5.1 | 9.2 |
| Haynes..... | 2874 | 18.3 | 13.8 | 9.0 | 0 | 0 | 14.1 | 5.1 | 8.7 |
| All stations: | | | | | | | | | |
| Kubanka..... | 1440 | 16.6 | 26.9 | 9.5 | 7.0 | 22.9 | 23.1 | 21.9 | 18.1 |
| Ghirka..... | 1517 | 11.4 | 23.3 | 15.0 | 7.1 | 18.8 | 22.5 | 16.8 | 16.4 |
| Rysting..... | 3022 | } 13.3 | 22.8 | 13.3 | 6.8 | 16.6 | 21.6 | 17.7 | 16.0 |
| Glyndon..... | 2873 | | | | | | | | |
| Haynes..... | 2874 | 12.8 | 23.1 | 10.7 | 5.9 | 12.2 | 21.1 | 14.1 | 14.9 |

^a Destroyed by hail in 1912.

^b Yield of Haynes Bluestem (C. I. No. 3021, Minn. No. 51).

^c Computed from the yields of the other varieties shown.

^d Results from State substation; work not cooperative. Edgeley numbers and C. I. numbers represent the same original stocks.

A second successive dry season occurred in 1911 at all stations except Moccasin. There a plentiful rainfall caused the production of large yields. Under the favorable conditions at Moccasin, the

¹ Salmon, Cecil. Sterile florets in wheat and other cereals. *In* Jour. Amer. Soc. Agron., v. 6, no. 1, p. 24-30, 2 pl. 1914.

Kubanka led in yield, the Rysting ranking second and Ghirka third. At the other stations, under severe drought conditions, the yields decreased from the north to the south, no yields being obtained at Highmore and Newell. At Williston and Dickinson the Ghirka again yielded considerably better than the other wheats, but did not retain this advantage at the stations farther south. Early rust, as well as drought, reduced its yield at Brookings. For all stations, however, the Ghirka ranked first, Kubanka second, Rysting-Glyndon third, and Haynes fourth.

The annual and seasonal precipitation at each station for the seven years is given in Table II. The averages of these data are included also in figure 3, to aid in the interpretation of the yield data.

TABLE II.—Annual and seasonal (April to July, inclusive) precipitation at seven experiment stations in three States of the northern Great Plains area, 1908 to 1914, inclusive.

| Station. | Precipitation (inches). | | | | | | | Average. |
|---------------------|-------------------------|-------|-------|-------|--------------------|-------|-------|--------------------|
| | 1908 | 1909 | 1910 | 1911 | 1912 | 1913 | 1914 | |
| Moccasin, Mont.: | | | | | | | | |
| Annual..... | 21.49 | 22.97 | 15.26 | 21.15 | ^a 15.00 | 14.96 | 15.67 | ^b 18.58 |
| Seasonal..... | 10.57 | 13.96 | 6.50 | 7.69 | ^a 7.93 | 9.32 | 9.38 | ^b 9.57 |
| Williston, N. Dak.: | | | | | | | | |
| Annual..... | 13.49 | 11.74 | 10.28 | 13.69 | 16.33 | 14.28 | 18.47 | 14.04 |
| Seasonal..... | 7.00 | 9.05 | 5.48 | 5.70 | 10.20 | 5.63 | 11.98 | 7.86 |
| Dickinson, N. Dak.: | | | | | | | | |
| Annual..... | 19.48 | 21.26 | 13.34 | 15.73 | ^a 19.06 | 11.93 | 22.74 | ^b 17.41 |
| Seasonal..... | 10.46 | 11.53 | 8.35 | 5.99 | ^a 12.46 | 5.31 | 16.79 | ^b 9.74 |
| Edgeley, N. Dak.: | | | | | | | | |
| Annual..... | 17.07 | 15.14 | 12.21 | 15.47 | 21.84 | 19.82 | 18.05 | 17.09 |
| Seasonal..... | 9.24 | 10.55 | 4.01 | 6.87 | 15.59 | 9.06 | 13.23 | 9.79 |
| Brookings, S. Dak.: | | | | | | | | |
| Annual..... | 32.34 | 22.34 | 12.78 | 24.95 | 23.18 | 16.58 | 24.15 | 22.33 |
| Seasonal..... | 19.60 | 10.64 | 6.74 | 10.62 | 14.95 | 10.69 | 14.09 | 12.47 |
| Highmore, S. Dak.: | | | | | | | | |
| Annual..... | 22.37 | 18.03 | 11.15 | 15.87 | 11.16 | 12.46 | 17.52 | 15.51 |
| Seasonal..... | 12.50 | 8.52 | 6.93 | 5.41 | 6.00 | 8.59 | 11.98 | 8.57 |
| Newell, S. Dak.: | | | | | | | | |
| Annual..... | 14.23 | 17.73 | 12.55 | 6.64 | 16.09 | 12.53 | 11.70 | 13.07 |
| Seasonal..... | 7.84 | 12.75 | 5.76 | 1.92 | 8.07 | 5.66 | 6.74 | 6.97 |
| All stations: | | | | | | | | |
| Annual..... | 20.07 | 18.46 | 12.51 | 16.21 | 17.72 | 14.65 | 18.33 | 17.15 |
| Seasonal..... | 11.03 | 11.00 | 6.25 | 6.31 | 10.96 | 7.75 | 12.03 | 9.28 |

^a These precipitation data are excluded from the average because the crop was destroyed by hail.

^b Average for only 6 years, excluding 1912.

In 1912 the varieties at Moccasin and Dickinson were destroyed by hail and no yields were obtained, although very favorable conditions existed until the hail occurred. At Williston, under similar favorable conditions, unusually large yields were obtained. At Edgeley and Brookings the conditions were fair, while at Highmore and Newell the third successive droughty year occurred. The Ghirka slightly outyielded the Kubanka at Williston and also led the other varieties at Highmore and Newell. At the latter station it was the only variety of the four that produced grain, thus showing again its drought-resisting ability. Kubanka produced a yield a third greater than the other varieties at both Edgeley and Brookings, due largely to the

greater rust resistance of the durum wheat. The rank in yield of the other three varieties at these two stations was reversed. For the five stations where results were obtained the Kubanka ranked first, Ghirka second, Rysting-Glyndon third, and Haynes fourth.

The favorable season of 1913 resulted in good yields at all stations except Highmore. The Kubanka led in yield at Moccasin, Williston, and Brookings, the Ghirka at Edgeley and Newell, the Rysting at Dickinson, and the Glyndon at Highmore. For all stations the Kubanka ranked first, Ghirka second, Rysting-Glyndon third, and Haynes fourth.

The season of 1914 was favorable at Moccasin and Williston, but drought, hail, and rust reduced the yields at the other stations. The yields have been computed for certain varieties which were not grown at three stations in 1914. The Haynes was discarded at Moccasin after 1912 and the Ghirka at Brookings after 1913. The Kubanka and Ghirka were unfortunately omitted, through error, from the test at Newell in 1914. The yields at Dickinson were damaged about two-thirds by hail. Rust reduced the yields at Edgeley, Brookings, and Highmore as much or more than drought.

The Kubanka produced the best yields at Williston, Edgeley, Brookings, and Highmore and equaled the yields of Ghirka and Rysting at Moccasin. The yield of Ghirka was relatively less at all stations except Moccasin than that usually obtained. For all stations the Kubanka ranked first, Rysting-Glyndon second, Ghirka third, and Haynes fourth.

SUMMARY OF YIELDS.

The averages for the seven years from 1908 to 1914, inclusive, covering a period of varying seasonal conditions, show a remarkable uniformity of results for the different varieties at each station. The Kubanka durum leads all varieties at all stations except Dickinson. Had the comparison included the results for 1907 at Dickinson the Kubanka would have outyielded the other varieties. A selection from Kubanka called Kubanka No. 8 (C. I. No. 4063) also outyielded the other varieties at Dickinson for the 5-year period from 1910 to 1914, inclusive.

The Ghirka has ranked second at Moccasin, Williston, Edgeley, and Newell, and tied with Rysting for second rank at Brookings. At Dickinson the Ghirka was first and at Highmore fourth.

The Rysting-Glyndon ranked third at Moccasin, Williston, Highmore, and Newell and tied at Brookings for second place. At Dickinson it ranked second and at Edgeley fourth.

The Haynes gave the lowest yields at Moccasin, Williston, Dickinson, Brookings, and Newell. It ranked third at Edgeley and second at Highmore.

The yield data presented are based on 182 actual determinations (crops destroyed by hail not considered). Yields have been computed in six other cases where the varieties were not sown and actual yields, therefore, were not obtained. This makes a total of 188 tests, or 47 tests for each of the four varieties. In three of these tests no yields of grain were obtained from all or from at least three of the varieties. In five other tests, either two or three of the varieties were equal in yield.

In 39 tests the four varieties can be definitely ranked. There are 24 possible combinations in which four varieties can rank. The varieties actually ranked in 15 of the possible 24 ways in the 39 tests. In annual yield, averaged for all stations, the varieties ranked in the same order only twice during the seven seasons. This shows conflicting annual results under varying seasonal conditions.

The varieties ranked first or equaled another variety for first rank, as follows: Kubanka 24 times, Ghirka 17 times, Rysting or Glyndon 6 times, and Haynes 2 times during 44 tests.

The average acre yield in 47 comparative tests at seven stations was 18.1 bushels for Kubanka, 16.4 bushels for Ghirka, 16 bushels for Rysting-Glyndon, and 14.9 bushels for Haynes. This shows a rather definite agreement of average results under varying seasonal conditions.

It is believed that the data presented fairly indicate what may be expected in yield from these four wheats in the northern Great Plains during a series of years.

The Ghirka variety has compared favorably with standard durum, Fife, and Bluestem varieties, yielding less than the durum but more than the Fife and Bluestem varieties on an average at seven stations during a 7-year period. While it is susceptible to rust in moist seasons, it has proved more drought resistant than the other wheats here studied, outyielding all others in the dry seasons.

MILLING AND BAKING QUALITY.

Quality as well as yield is an important consideration in determining the value of a variety. Quality of wheat is thought of in many different ways. To the farmer it means grade; to the miller and baker it is the ability of a wheat to produce a high percentage of flour and a loaf of large volume. Relatively few data have been gathered on the quality of the Ghirka wheat as compared with those on yield.

Hard spring common wheats grade as No. 1 Hard, No. 1 Northern, No. 2 Northern, etc. The Ghirka wheat when marketed has been included in the Northern grade. The somewhat soft kernel prevents its ever grading as No. 1 Hard. No extensive tests have been made of the crushing point of the Ghirka kernel compared with

that of other varieties. A preliminary test, however, indicates the Ghirka to be about three-fourths as hard as the Haynes Bluestem and about one-half as hard as Kubanka durum wheat. This confirms the general opinion that Ghirka is a semi-hard rather than a hard wheat.

Many milling and baking tests have been made of Ghirka wheat grown at various stations, but the methods used by the investigators have varied. For several years the Office of Cereal Investigations and the North Dakota Agricultural Experiment Station have studied the milling and baking qualities of the principal wheats grown at the Dickinson substation. During three years, 1911, 1913, and 1914, the method used was the same. These data compare the milling and baking qualities of the Ghirka with those of the wheats already compared with it in regard to yield. The data for the Ghirka, Kubanka, and Haynes varieties are strictly comparable. The results from the Rysting in 1911 and 1913 have been combined with those from the Glyndon in 1914 to make the data somewhat comparable with those on yield. The annual and average results are given in Table III. In figure 4 only the average results are shown.




FIG. 4.—Diagram showing the average results of milling and baking tests of four varieties of wheat grown at the Dickinson substation during 1911, 1913, and 1914.

and the color of the loaf was better than that from the Kubanka. The Haynes gave the highest yield of straight flour, as well as the largest volume of loaf. The Rysting had the best color and texture of loaf.

In 1913, data on the crude-protein content of the kernel and on loss in milling were obtained for the first time. In that year the Ghirka was highest in loaf volume but lowest in crude-protein con-

tent and yield of straight flour, and it showed the greatest loss in milling. The low yield of flour and the loss in milling are especially pronounced. This is partly due to a peculiar fluffiness of the Ghirka flour. The Haynes gave the highest yield of straight flour and tied Kubanka for the highest score in texture of loaf. The Rysting had the highest percentage of crude protein in the wheat and the best color of loaf.

The data for 1914 show that the Ghirka is the poorest in crude-protein content and color and texture of loaf and shows the greatest loss in milling. The Haynes leads in crude-protein content and texture of loaf. The Glyndon yields more straight flour, produces the best volume of loaf, and ties with the Haynes in color of loaf.

The 3-year average of these data shows that the Ghirka variety has not compared favorably in quality with standard Bluestem, Fife, and durum wheats. It ranks relatively low in market grade, in crude protein, color and texture of loaf, and yield of straight flour, and has a large loss in milling. The volume of loaf is usually good, although averaging less than that of Bluestem.

TABLE III.—*Milling and baking tests of Ghirka and three other varieties of wheat grown at the Dickinson (N. Dak.) substation in 1911, 1913, and 1914.^a*

| Variety. | C. I. No. | 1911 | 1913 | 1914 | Average. |
|---|-----------|-------|-------|-------|----------|
| Crude protein (N × 5.7) in wheat (per cent): | | | | | |
| Kubanka..... | 1440 | | 15.50 | 14.99 | 15.25 |
| Ghirka..... | 1517 | | 14.36 | 13.52 | 13.94 |
| Rysting..... | b 3022 | | 15.80 | 15.33 | 15.57 |
| Glyndon..... | b 2873 | | | | |
| Haynes..... | 2874 | | 14.99 | 15.50 | 15.25 |
| Yield of straight flour (per cent): | | | | | |
| Kubanka..... | 1440 | 71.2 | 69.7 | 66.9 | 69.3 |
| Ghirka..... | 1517 | 68.7 | 65.3 | 67.3 | 67.1 |
| Rysting..... | b 3022 | 71.2 | 70.6 | 68.4 | 70.1 |
| Glyndon..... | b 2873 | | | | |
| Haynes..... | 2874 | 73.5 | 71.2 | 66.6 | 70.4 |
| Loss in milling (per cent): | | | | | |
| Kubanka..... | 1440 | | 2.57 | .82 | 1.70 |
| Ghirka..... | 1517 | | 5.94 | 2.80 | 4.37 |
| Rysting..... | b 3022 | | 3.22 | 1.80 | 2.51 |
| Glyndon..... | b 2873 | | | | |
| Haynes..... | 2874 | | 2.68 | 2.37 | 2.53 |
| Volume of loaf (cubic centimeters): | | | | | |
| Kubanka..... | 1440 | 1,970 | 1,985 | 2,435 | 2,130 |
| Ghirka..... | 1517 | 2,150 | 2,270 | 2,648 | 2,356 |
| Rysting..... | b 3022 | 2,060 | 2,000 | 2,945 | 2,335 |
| Glyndon..... | b 2873 | | | | |
| Haynes..... | 2874 | 2,320 | 2,260 | 2,900 | 2,493 |
| Color of loaf (per cent): | | | | | |
| Kubanka..... | 1440 | 83 | 93 | 96.0 | 90.7 |
| Ghirka..... | 1517 | 85 | 96 | 91.5 | 90.8 |
| Rysting..... | b 3022 | 97 | 99 | 97.0 | 97.7 |
| Glyndon..... | b 2873 | | | | |
| Haynes..... | 2874 | 95 | 96 | 97.0 | 96.0 |
| Texture of loaf (per cent): | | | | | |
| Kubanka..... | 1440 | 86 | 94 | 95.0 | 91.7 |
| Ghirka..... | 1517 | 85 | 93 | 93.0 | 90.3 |
| Rysting..... | b 3022 | 87 | 90 | 95.5 | 90.8 |
| Glyndon..... | b 2873 | | | | |
| Haynes..... | 2874 | 86 | 94 | 96.0 | 92.0 |

^a The grain was milled and the bread baked at the North Dakota Agricultural Experiment Station.

^b Results for 1911 and 1913 are from Rysting (C. I. No. 3022); for 1914, from Glyndon (C. I. No. 2873).

The Haynes has the highest average yield of flour, volume of loaf, and texture of loaf. The Rysting-Glyndon has the highest average crude-protein content and color of loaf. The Kubanka shows the smallest loss in milling.

The data given show that the Haynes is superior to the Ghirka in all of the six characters studied. The Rysting-Glyndon is superior in all characters except loaf volume, and the Kubanka is superior in all characters except volume and color of loaf.

IMPROVEMENT BY SELECTION.

The high-yielding and drought-resistant power of the Ghirka wheat was early recognized at the Dickinson substation. A further improvement in the yield and quality of the variety was sought through isolating a large number of pure lines and comparing them with the



FIG. 5.—Pure-line selections of Ghirka Spring wheat growing in eightieth-acre plats at the Dickinson substation in 1913.

original. This work was started in 1909, when about 300 heads were selected from a plat of the Ghirka wheat.¹ The kernels from these were sown in head rows in 1910. Yield data were obtained, and all the best, a large number of the medium, and some of the poorest were selected for sowing in 1911. Of these selections, 104 were sown in 17-foot rows for comparative yields, with the original Ghirka mass variety used as a check every third row. Eighty of the selections were sown also in rows of varying length to increase

¹ The experiments to improve the Ghirka Spring variety by pure-line selection were suggested by Supt. L. R. Waldron, of the Dickinson substation, who has maintained his interest in the work throughout the entire period. The original selections were made by Mr. Charles H. Clark, who was in charge of the cereal work at the Dickinson substation in 1909 and 1910. From 1911 to July 25, 1914, the writer was in direct charge of the cooperative cereal experiments at Dickinson. On the latter date he was succeeded by Mr. Ralph W. Smith, who continues in charge of the work as Scientific Assistant in Cereal Investigations.

the seed. Considerable variation was observed in the different pure lines, not only in respect to yield but also in the color and texture of the kernel, morphological characters of the plant, etc.

In 1912 the same 104 selections were sown in duplicated 17-foot rows with checks. About 20 were sown also in hundredth-acre plats. These tests were all destroyed by hail.




FIG. 6.—Diagram showing the annual and average actual and computed yields of Ghirka Spring wheat (C. I. No. 1517) used as a check and 30 pure lines selected therefrom, grown at the Dickinson substation in 1913 and 1914.

In 1913 the entire 104 selections were again sown in 17-foot rows from reserve stocks of seed. In addition, 86 of them were grown in 108-foot rows and 24 in plats (fig. 5) of one-eightieth acre. This series of tests gave considerable information on the behavior of these pure lines.

Samples of the grain from the 24 pure lines grown in field plats were used in milling and baking tests in cooperation with the North Dakota Agricultural Experiment Station.

In 1914 the work was continued. Besides the nursery work, 24 selections were again grown in field plats. Eighteen of these were the same as those grown in plats in 1913. The wheats were greatly damaged by hail which fell shortly before the crop was ripe. The plats were harvested, however, and yield data obtained. Milling and baking tests were also made on a number of the selections.

The annual agronomic data for the pure lines grown in plats in 1913 and 1914 are given in Table IV, together with data from the check plats of the original Ghirka, C. I. No. 1517. In figure 6 are shown the results obtained in 1913 and 1914 and the average for the two years. The pure lines are arranged in the actual order of the sowings, and the actual and computed yields are shown separately. The actual yield of the check plats and the average for all checks are shown by different lines.

Improvement in the quality of Ghirka wheat is more important than increase in yield. It was hoped that a pure line would be found which had the combined characters necessary for good quality and high yield. A study of the milling and baking data obtained during the two years 1913 and 1914 on the highest yielding selections indicates that the desired result was obtained. These data are shown in Table V.

Figure 6 shows that in 1913 the Ghirka selections Nos. 4, 5, 17, 31, 66, and 72 were outstanding in regard to yield and that many others were better than the parent mass variety. The data for 1914 show




FIG. 7.—Diagram showing the average results of milling and baking tests of Ghirka Spring wheat (C. I. No. 1517) and four pure lines selected therefrom, grown at the Dickinson substation in 1913 and 1914.

that Nos. 4, 5, 50, 66, and 72 produced outstanding yields. An average of the yields of the pure lines grown both years shows Nos. 4, 5, 66, and 72 to be the best four selections when both actual and computed yields are considered, each showing an increase of more than

3 bushels over the parent variety. No definite conclusions can be drawn from the results of only two years, but the data indicate that improvement in yield is possible and that progress is being made. During the years covered by these data rust was not prevalent and no data were obtained on the rust-resisting ability of the different selections.

The average data for the four highest yielding pure lines and the parent mass variety are shown graphically in figure 7. The data given for the pure lines each year are an average of one milling and two bakings and for the parent variety are an average of these data from two plats, or an average of two millings and four bakings.

In 1913, Ghirka selections Nos. 4 and 66 exceeded the original Ghirka in crude-protein content and in yield of straight flour, gave a smaller loss in milling, and produced loaves of greater volume, which also scored higher in color and texture. This was an improvement in all of the characters here studied. Nos. 5 and 72 exceeded the original in some characters but failed to equal it in others.

In 1914, No. 4 gave the highest volume of loaf, but was low in crude protein and yield of straight flour, had a high loss in milling, and scored low in color. No. 66 scored low in volume of loaf and in texture, but otherwise exceeded the check. For the second year it showed an unusually low loss in milling.

Selection No. 5 led in crude protein and yield of straight flour, was superior to the original in volume and texture of loaf, and lost less in milling. Selection No. 72 was low in yield of flour and in volume, color, and texture of loaf. None of the four highest yielding pure lines exceeded the original variety in all characters in 1914.

The average for the two years shows that selection No. 4 was superior to the unselected Ghirka in all characters except crude-protein content and No. 66 in all characters except volume of loaf. No. 5 shows a decrease in crude protein and in color of loaf, while No. 72 shows a decrease in the volume, color, and texture of loaf. While none of the four best yielding selections has exceeded the mass variety in all of the characters studied, a gain in some of the characters offsets the loss in others, and the data thus far obtained indicate that at least the first three selections mentioned are superior in quality to the parent mass variety.

TABLE IV.—Agronomic data for unselected Ghirka wheat, used as a check, and 30 pure lines selected therefrom, grown at the Dickinson (N. Dak.) substation in 1913 and 1914.

| Ghirka and selections. | C. I. No. | Date. | | Planting to maturity. | Stand. | | Height. | Yield per acre. | | | | Weight per bushel. | |
|------------------------------|-----------|---------------|-------------|-----------------------|------------|------------------|---------|-----------------|---------|--------|-----------|--------------------|--------|
| | | Headed, July— | Ripe, Aug.— | | Days. | Plants per acre. | | Estimated. | Actual. | | Computed. | | |
| | | | | | | | | | Grain. | Straw. | Grain. | | Straw. |
| Season of 1913: ¹ | | | | | Thousands. | Per ct. | In. | Bu. | Cwt. | Bu. | Cwt. | Lbs. | |
| Ghirka..... | 1517 | 14 | 12 | 113 | 640 | 86 | | 27.3 | 20.4 | 30.0 | 24.6 | 61 | |
| Selections— | | | | | | | | | | | | | |
| No. 4..... | 4413 | 14 | 12 | 113 | 696 | 94 | 32.1 | 35.6 | 25.3 | 39.0 | 31.0 | 62 | |
| No. 5..... | 4414 | 12 | 9 | 110 | 776 | 105 | 34.0 | 40.5 | 23.3 | 44.3 | 28.8 | 62.5 | |
| No. 12..... | 4415 | 12 | 12 | 113 | 688 | 93 | 33.4 | 35.3 | 22.4 | 38.7 | 28.0 | 62.5 | |
| No. 13..... | | 10 | 10 | 111 | 844 | 114 | 34.2 | 31.3 | 19.6 | 32.3 | 25.2 | 61.5 | |
| Ghirka..... | 1517 | 12 | 12 | 113 | 700 | 95 | | 27.3 | 19.2 | 30.0 | 24.6 | 62.3 | |
| Selections— | | | | | | | | | | | | | |
| No. 15..... | 4416 | 16 | 15 | 116 | 560 | 76 | 33.2 | 28.8 | 19.8 | 31.3 | 24.3 | 61.5 | |
| No. 17..... | 4417 | 14 | 12 | 113 | 724 | 98 | 34.5 | 35.2 | 21.7 | 36.1 | 25.5 | 62.5 | |
| No. 19..... | | 12 | 12 | 113 | 860 | 116 | 34.3 | 27.7 | 20.2 | 28.9 | 22.9 | 61.5 | |
| No. 20..... | | 10 | 9 | 110 | 692 | 94 | 32.3 | 32.2 | 23.5 | 33.2 | 25.6 | 61 | |
| Ghirka..... | 1517 | 12 | 12 | 113 | 700 | 95 | | 29.5 | 23.5 | 30.0 | 24.6 | 62 | |
| Selections— | | | | | | | | | | | | | |
| No. 21..... | | 12 | 12 | 113 | 608 | 82 | 32.4 | 33.0 | 24.3 | 31.8 | 24.1 | 62 | |
| No. 23..... | 4418 | 14 | 12 | 113 | 868 | 117 | 35.0 | 34.7 | 25.6 | 31.9 | 24.2 | 62.5 | |
| No. 24..... | 4419 | 14 | 12 | 113 | 728 | 98 | 34.0 | 36.0 | 25.2 | 31.6 | 22.6 | 61.5 | |
| No. 25..... | 4420 | 14 | 12 | 113 | 840 | 113 | 34.3 | 36.2 | 26.9 | 30.4 | 22.2 | 62 | |
| Ghirka..... | 1517 | 12 | 12 | 113 | 680 | 92 | | 37.2 | 30.1 | 30.0 | 24.6 | 62.5 | |
| Selections— | | | | | | | | | | | | | |
| No. 29..... | 4421 | 10 | 9 | 110 | 860 | 116 | 32.3 | 33.2 | 25.7 | 27.3 | 21.5 | 61 | |
| No. 31..... | | 12 | 12 | 113 | 768 | 104 | 32.7 | 38.0 | 29.2 | 31.9 | 25.0 | 62 | |
| No. 36..... | | 12 | 12 | 113 | 820 | 111 | 32.3 | 34.1 | 28.0 | 29.3 | 24.5 | 62.5 | |
| No. 37..... | | 12 | 12 | 113 | 836 | 113 | 31.8 | 33.3 | 26.0 | 29.3 | 23.3 | 62 | |
| Ghirka..... | 1517 | 12 | 12 | 113 | 648 | 88 | | 33.3 | 26.8 | 30.0 | 24.6 | 62 | |
| Selections— | | | | | | | | | | | | | |
| No. 40..... | 4422 | 12 | 12 | 113 | 768 | 104 | 31.7 | 32.0 | 23.2 | 29.7 | 21.2 | 62 | |
| No. 47..... | | 12 | 13 | 114 | 752 | 102 | 30.4 | 28.8 | 27.9 | 27.6 | 25.4 | 61 | |
| No. 61..... | | 12 | 9 | 110 | 688 | 93 | 34.3 | 31.7 | 26.7 | 31.4 | 24.1 | 62 | |
| No. 65..... | 4424 | 14 | 9 | 110 | 896 | 121 | 33.4 | 33.0 | 25.9 | 33.8 | 23.4 | 61.5 | |
| No. 66..... | 4425 | 12 | 12 | 113 | 828 | 112 | 31.6 | 35.2 | 25.9 | 37.3 | 23.1 | 61.5 | |
| No. 68..... | 4426 | 10 | 10 | 111 | 792 | 107 | | 31.3 | 22.0 | 34.5 | 19.7 | 60.5 | |
| No. 72..... | 4427 | 11 | 12 | 113 | 720 | 97 | 33.9 | 32.0 | 25.6 | 36.6 | 22.9 | 62.5 | |
| Ghirka..... | 1517 | 12 | 12 | 113 | 684 | 92 | | 25.2 | 27.7 | 30.0 | 24.6 | 61 | |
| Selections— | | | | | | | | | | | | | |
| No. 74..... | | 12 | 12 | 113 | 524 | 71 | 34.2 | 19.3 | 17.6 | 24.0 | 22.8 | 61 | |
| Season of 1914: ² | | | | | | | | | | | | | |
| Ghirka..... | 1517 | 10 | 8 | 112 | 616 | 114 | 41.5 | 6.4 | 17.3 | 10.3 | 19.0 | 51 | |
| Selections— | | | | | | | | | | | | | |
| No. 4..... | 4413 | 10 | 8 | 112 | 532 | 99 | | 11.2 | 20.2 | 15.8 | 21.6 | 52.5 | |
| No. 5..... | 4414 | 10 | 8 | 112 | 454 | 84 | 37.2 | 13.2 | 25.7 | 16.4 | 26.9 | 52.5 | |
| No. 12..... | 4415 | 10 | 8 | 112 | 436 | 81 | 39.8 | 7.6 | 9.8 | 8.5 | 10.1 | 53 | |
| No. 17..... | 4417 | 10 | 8 | 112 | 490 | 91 | 39.2 | 8.8 | 9.1 | 9.0 | 9.1 | 51.5 | |
| No. 20..... | | 10 | 8 | 112 | 460 | 85 | 36.7 | 6.4 | 24.0 | 5.9 | 23.6 | 49 | |
| Ghirka..... | 1517 | 10 | 8 | 112 | 596 | 110 | 41.5 | 12.0 | 19.7 | 10.3 | 19.0 | | |
| Selections— | | | | | | | | | | | | | |
| No. 21..... | | 12 | 8 | 112 | 460 | 85 | 39.9 | 10.4 | 20.6 | 8.9 | 20.1 | 53 | |
| No. 24..... | 4419 | 10 | 8 | 112 | 464 | 86 | 39.9 | 10.0 | 23.8 | 8.6 | 23.2 | 51.5 | |
| No. 25..... | 4420 | 10 | 8 | 112 | 440 | 82 | 41.1 | 9.2 | 20.4 | 7.2 | 20.2 | 52 | |
| No. 29..... | 4421 | 8 | 8 | 112 | 456 | 85 | 39.2 | 10.0 | 19.0 | 8.6 | 18.9 | 49 | |
| No. 31..... | | 10 | 8 | 112 | 472 | 87 | 39.4 | 9.2 | 17.5 | 7.2 | 17.6 | 52 | |
| Ghirka..... | 1517 | 10 | 8 | 112 | 644 | 119 | 39.8 | 12.0 | 18.7 | 10.3 | 19.0 | 52 | |
| Selections— | | | | | | | | | | | | | |
| No. 36..... | | 10 | 8 | 112 | 552 | 102 | 41.9 | 11.2 | 22.1 | 9.9 | 22.1 | 51.5 | |
| No. 40..... | 4422 | 11 | 8 | 112 | 472 | 87 | 40.2 | 10.4 | 28.3 | 9.6 | 28.0 | 52 | |
| Selections— | | | | | | | | | | | | | |
| No. 47..... | | 10 | 8 | 112 | 644 | 119 | 38.1 | 9.6 | 15.4 | 9.2 | 15.0 | 51.5 | |
| No. 50..... | 4423 | 9 | 8 | 112 | 636 | 118 | 40.3 | 12.8 | 25.0 | 12.7 | 24.1 | 52 | |
| No. 55..... | | 9 | 8 | 112 | 580 | 108 | 41.3 | 10.4 | 23.5 | 10.7 | 22.4 | 53.5 | |
| Ghirka..... | 1517 | 10 | 8 | 112 | 604 | 112 | 40.4 | 9.6 | 20.2 | 10.3 | 19.0 | 51.5 | |
| Selections— | | | | | | | | | | | | | |
| No. 57..... | | 9 | 8 | 112 | 628 | 116 | 43.5 | 11.2 | 19.2 | 11.9 | 18.7 | 52 | |
| No. 59..... | | 11 | 8 | 112 | 532 | 99 | 38.8 | 10.0 | 27.6 | 10.4 | 27.9 | 51 | |
| No. 61..... | | 10 | 8 | 112 | 592 | 110 | 40.9 | 10.0 | 17.0 | 10.3 | 18.0 | 53 | |
| No. 62..... | | 10 | 8 | 112 | 516 | 96 | 40.7 | 9.2 | 22.3 | 9.4 | 24.5 | 50.5 | |
| No. 65..... | 4424 | 10 | 8 | 112 | 544 | 101 | 37.1 | 11.2 | 20.2 | 11.2 | 23.1 | 51 | |
| Ghirka..... | 1517 | 10 | 8 | 112 | 512 | 95 | 41.6 | 10.4 | 15.8 | 10.3 | 19.0 | 52 | |
| Selections— | | | | | | | | | | | | | |
| Nos. 4 and 66..... | | 9 | 8 | 112 | 528 | 98 | 40.1 | 10.0 | 21.8 | 9.8 | 24.5 | 52.5 | |
| No. 66..... | 4425 | 9 | 8 | 112 | 584 | 108 | 42.6 | 12.4 | 27.1 | 12.0 | 28.8 | 52.5 | |
| No. 68..... | 4426 | 10 | 8 | 112 | 524 | 97 | 39.2 | 9.6 | 18.2 | 9.2 | 18.2 | 51 | |
| No. 72..... | 4427 | 10 | 8 | 112 | 496 | 92 | | 14.4 | 29.8 | 13.6 | 28.2 | 51.5 | |
| No. 102..... | | 14 | 8 | 112 | 664 | 123 | | 8.4 | 24.7 | 7.8 | 22.3 | 49 | |
| Ghirka..... | 1517 | 11 | 8 | 112 | 596 | 110 | 42.4 | 11.2 | 22.1 | 10.3 | 19.0 | 52.5 | |

¹ Sown Apr. 21, emerged May 10, harvested Aug. 15, 1913.² Sown Apr. 18, emerged May 2, harvested Aug. 10, 1914.

TABLE V.—*Milling and baking data for Ghirka wheat, used as a check, and 25 pure lines selected therefrom, grown at the Dickinson (N. Dak.) substation in 1913 and 1914.*

| Plat No. | Variety. | C. I. No. | Crude protein in wheat (N×5.7). | Yield of flour. | Loss in mill-ing. | Loaf. | | |
|-----------------|-------------|-----------|---------------------------------|-----------------|-------------------|----------|--------|-----------|
| | | | | | | Vol-ume. | Color. | Tex-ture. |
| Season of 1913: | | | | | | | | |
| Selections— | | | | | | | | |
| 2 | No. 4..... | 4413 | 14.36 | 71.2 | 3.35 | 2,450 | 96 | 94 |
| 3 | No. 5..... | 4414 | 13.45 | 69.6 | 5.65 | 2,300 | 94 | 93 |
| 4 | No. 12..... | 4415 | 14.42 | 68.7 | 4.75 | 2,320 | 92 | 94 |
| 5 | No. 13..... | | 14.99 | 67.1 | 5.78 | 2,130 | 84 | 88 |
| 7 | No. 15..... | 4416 | 15.28 | 69.9 | 7.39 | 1,860 | 92 | 88 |
| 8 | No. 17..... | 4417 | 13.97 | 68.5 | 5.14 | 2,200 | 99 | 92 |
| 9 | No. 19..... | | 15.28 | 69.1 | 4.50 | 2,265 | 96 | 93 |
| 10 | No. 20..... | | 16.07 | 68.9 | 5.10 | 1,885 | 90 | 90 |
| 12 | No. 21..... | | 14.82 | 70.5 | 2.85 | 2,155 | 94 | 92 |
| 13 | No. 23..... | 4418 | 14.36 | 69.8 | 4.50 | 2,150 | 89 | 90 |
| 14 | No. 24..... | 4419 | 14.54 | 67.2 | 5.98 | 2,350 | 89 | 94 |
| 15 | No. 25..... | 4420 | 15.16 | 70.3 | 4.01 | 2,065 | 89 | 90 |
| 17 | No. 29..... | 4421 | 15.62 | 67.5 | 7.10 | 1,790 | 94 | 86 |
| 18 | No. 31..... | | 15.28 | 70.1 | 4.17 | 1,985 | 90 | 92 |
| 19 | No. 36..... | | 15.16 | 69.8 | 4.32 | 2,030 | 91 | 90 |
| 20 | No. 37..... | | 14.54 | 70.4 | 4.57 | 1,895 | 86 | 90 |
| 22 | No. 40..... | 4422 | 14.54 | 72.0 | 3.10 | 2,120 | 95 | 93.5 |
| 23 | No. 47..... | | 14.88 | 70.7 | 4.77 | 2,110 | 96 | 93 |
| 24 | No. 61..... | | 14.71 | 69.2 | 5.73 | 2,080 | 93 | 93 |
| 25 | No. 65..... | 4424 | 14.48 | 68.9 | 5.17 | 2,050 | 94 | 94 |
| 26 | No. 66..... | 4425 | 14.76 | 72.4 | 1.46 | 2,097 | 95 | 93.5 |
| 27 | No. 68..... | 4426 | 15.50 | 67.2 | 7.68 | 1,825 | 89 | 85 |
| 28 | No. 72..... | 4427 | 14.99 | 72.2 | 3.49 | 1,975 | 92 | 92 |
| 30 | No. 74..... | | 14.71 | 71.7 | 4.21 | 1,895 | 90 | 89 |
| 21 | Ghirka..... | 1517 | 13.88 | 69.6 | 5.51 | 2,011 | 93 | 91.5 |
| 29 | | | | | | | | |
| Season of 1914: | | | | | | | | |
| Selections— | | | | | | | | |
| 2 | No. 4..... | 4413 | 12.94 | 67.2 | 3.06 | 2,685 | 90 | 95 |
| 3 | No. 5..... | 4414 | 14.54 | 69.0 | 1.39 | 2,660 | 89 | 94 |
| 9 | No. 24..... | 4419 | 13.22 | 66.9 | 2.67 | 2,675 | 94 | 92 |
| 11 | No. 29..... | 4421 | 13.97 | 66.0 | 2.06 | 2,590 | 90 | 92 |
| 15 | No. 40..... | 4422 | 14.36 | 65.7 | 2.60 | 2,320 | 93 | 89 |
| 17 | No. 50..... | 4423 | 14.71 | 68.0 | 1.80 | 2,405 | 93 | 90 |
| 27 | No. 66..... | 4425 | 14.36 | 67.9 | .80 | 2,520 | 93 | 92.5 |
| 29 | No. 72..... | 4427 | 14.19 | 66.5 | 1.73 | 2,470 | 91 | 90.5 |
| 13 | Ghirka..... | 1517 | 14.02 | 67.3 | 2.80 | 2,648 | 91.5 | 93 |
| 25 | | | | | | | | |

A comparison of these three pure lines with standard varieties will serve to show their relative values. Five such varieties have been grown and a comparison is made of their annual and average yields per acre, percentages of straight flour, and volumes of loaf in Table VI. While the data on yield are not strictly comparable, owing to the difference noted in the size of plats, they show that the selections possess the ability to produce comparatively high yields. In percentage of straight flour the average shows that selection No. 66 exceeds any of the other varieties studied, while selections Nos. 5 and 4 exceed the Haynes, Marquis, and Red Fife. In volume of loaf the Haynes slightly exceeds selection No. 4, which, in turn, is superior to the Red Fife, Preston, Marquis, and Kubanka, while selection No. 5 is superior to the Preston, Marquis, and Kubanka, and selection No. 66 is superior to the Kubanka.

These selections and several others are being further tested to determine more accurately their true value before any are distributed.

TABLE VI.—Yields of grain and of flour and volumes of loaf obtained from three pure-line selections of Ghirka and five standard varieties of hard spring wheat grown at the Dickinson (N. Dak.) substation in 1913 and 1914.

[The varieties are arranged in the order of their average yields.]

| Variety. | C. I. No. | 1913 | 1914 | Average. |
|-------------------------------------|-----------|--------|--------|----------|
| Yield per acre (bushels): | | | | |
| Ghirka No. 5..... | 4414 | a 40.5 | b 13.2 | 26.9 |
| Ghirka No. 66..... | 4425 | a 35.2 | b 12.4 | 23.8 |
| Ghirka No. 4..... | 4413 | a 35.7 | b 11.2 | 23.5 |
| Kubanka..... | 1440 | c 31.2 | d 14.2 | 22.7 |
| Red Fife..... | 3329 | c 28.3 | d 10.2 | 19.3 |
| Preston..... | 3081 | c 25.6 | d 12.9 | 19.3 |
| Marquis..... | 3641 | c 24.0 | d 14.0 | 19.0 |
| Haynes..... | 2874 | c 24.8 | d 8.3 | 16.6 |
| Yield of straight flour (per cent): | | | | |
| Ghirka No. 66..... | 4425 | 72.4 | 67.9 | 70.2 |
| Kubanka..... | 1440 | 71.8 | 66.9 | 69.4 |
| Preston..... | 3081 | 70.7 | 68.1 | 69.4 |
| Ghirka No. 5..... | 4414 | 69.6 | 69.0 | 69.3 |
| Ghirka No. 4..... | 4413 | 71.2 | 67.2 | 69.2 |
| Haynes..... | 2874 | 71.2 | 66.6 | 68.9 |
| Marquis..... | 3641 | 69.2 | 63.8 | 66.5 |
| Red Fife..... | 3329 | 71.8 | 60.0 | 65.9 |
| Volume of loaf (cubic centimeters): | | | | |
| Haynes..... | 2874 | 2,260 | 2,900 | 2,580 |
| Ghirka No. 4..... | 4413 | 2,450 | 2,685 | 2,568 |
| Red Fife..... | 3329 | 2,220 | 2,775 | 2,498 |
| Ghirka No. 5..... | 4414 | 2,300 | 2,660 | 2,480 |
| Preston..... | 3081 | 2,170 | 2,785 | 2,478 |
| Marquis..... | 3641 | 2,250 | 2,475 | 2,363 |
| Ghirka No. 66..... | 4425 | 2,097 | 2,520 | 2,309 |
| Kubanka..... | 1440 | 1,985 | 2,435 | 2,210 |

a Plats one-eightieth of an acre.

c Plats one-tenth of an acre.

b Plats one forty-eighth of an acre.

d Plats one forty-eighth of an acre, replicated four times.

It is possible that a mixture of two or three of the best selections may result in a further improvement in yield or quality. A preliminary test to determine this, combining selections Nos. 4 and 66, is already under way. Crosses have been made between these best pure lines and other varieties of hard spring common wheat, and promising selections of the progeny of the second and fourth generations are now in existence. From these a still further improvement in quality and in rust resistance is expected, while retaining the high-yielding and drought-resistant qualities of the Ghirka wheat.

CONCLUSIONS.

Ghirka Spring wheat, a variety of commercial importance in Russia, has proved adapted to the northern part of the Great Plains area of the United States. The variety is susceptible to rust in moist seasons and in humid areas, but it has proved to be a valuable drought-resistant wheat.

Tests at seven experiment stations in the northern Plains area, covering a period of seven years, have shown that on an average the Ghirka Spring has yielded more than the Rysting Fife and Haynes Bluestem common wheats, but less than the Kubanka durum wheat. The quality of the Ghirka, however, is inferior to that of these standard wheats.

Experiments are being made at the Dickinson (N. Dak.) substation to improve both the yield and the quality of the Ghirka wheat by isolating superior pure lines. Many pure lines have been tested, and among them selections Nos. 4, 5, and 66 have thus far proved superior to the others and to the original mass variety in both yield and quality. They also compare favorably with the standard spring wheats of the Great Plains area in quality as well as in yield. The selections are also proving valuable material for crossing with varieties possessing greater rust resistance and high quality of grain.

From the data given it is shown that improvement in yield and quality is possible from pure-line selection and that good results are being obtained.

**PUBLICATIONS OF THE UNITED STATES DEPARTMENT OF AGRICULTURE
RELATING TO THE IMPROVEMENT OF WHEAT.**

AVAILABLE FOR FREE DISTRIBUTION.

- Cereal Experiments at Dickinson, N. Dak. (Department Bulletin 33.)
Cereal Experiments at the Williston Substation. (Department Bulletin 270.)
Cereal Experiments on the Belle Fourche Experiment Farm. (Department Bulletin 297.)
Cereal Experiments in Maryland and Virginia. (Department Bulletin 336.)
Alaska and Stoner, or "Miracle," Wheats: Two Varieties Much Misrepresented. (Department Bulletin 357.)
Cereal Experiments at the Judith Basin Substation, Moccasin, Mont. (Department Bulletin 398.)
Experiments with Marquis Wheat. (Department Bulletin 400.)
Cereal Experiments at the Akron Field Station, Akron, Colo. (Department Bulletin 402.)
Cereal Experiments on the Cheyenne Experiment Farm, Archer, Wyo. (Department Bulletin 430.)
The Smuts of Wheat, Oats, Barley, and Corn. (Farmers' Bulletin 507.)
Durum Wheat. (Farmers' Bulletin 534.)
Culture of Winter Wheat in the Eastern United States. (Farmers' Bulletin 596.)
Winter Wheat Varieties for the Eastern United States. (Farmers' Bulletin 616.)
Growing Hard Spring Wheat. (Farmers' Bulletin 678.)
Varieties of Hard Spring Wheat. (Farmers' Bulletin 680.)
Marquis Wheat. (Farmers' Bulletin 732.)
Hard Wheats Winning Their Way. (Separate 649 from Yearbook 1916.)

**FOR SALE BY THE SUPERINTENDENT OF DOCUMENTS, GOVERNMENT PRINTING
OFFICE, WASHINGTON, D. C.**

- Cereal Investigations at the Nephi Substation. (Department Bulletin 30.) Price, 10 cents.
Experiments with Wheat, Oats, and Barley in South Dakota. (Department Bulletin 39.) Price, 10 cents.
Spring Wheat in the Great Plains Area: Relation of Cultural Methods to Production. (Department Bulletin 214.) Price, 10 cents.
The Commercial Status of Durum Wheat. (Bureau of Plant Industry Bulletin 70.) Price, 10 cents.
Improving the Quality of Wheat. (Bureau of Plant Industry Bulletin 78.) Price, 10 cents.
Improvement of the Wheat Crop in California. (Bureau of Plant Industry Bulletin 178.) Price, 10 cents.
Winter Wheat in Western South Dakota. (Bureau of Plant Industry Circular 79.) Price, 5 cents.

